

AD-A084 837

CALIFORNIA UNIV LOS ANGELES DEPT OF CHEMISTRY

F/8 7/3

REACTION OF (IR(CYCLO-OCTA-1,5-DIENE) (P(P-TOLYL)3)2) (NID0-7,8---ETC(U)

MAY 80 J A DOI, R G TELLER, M F HANTHORNE

N00014-76-C-0390

UNCLASSIFIED

TR-109

NL

/ (P-)  
AD  
A084 837



END  
DATE  
FILMED  
6-80  
DTIC

ADA084837

62  
**LEVEL**

**(12)**

OFFICE OF NAVAL RESEARCH

15  
Contract No. N00014-76-C-0390

Task No. NR 053-608

TECHNICAL REPORT NO. 109

*1 Summary*  
**Reaction of [Ir(cyclo-octa-1,5-diene){P(p-tolyl)}<sub>3</sub>]<sub>2</sub>**

**TR-107**

**[nido-7,8-C<sub>2</sub>B<sub>9</sub>H<sub>12</sub>] with Hydrogen; Synthesis and X-Ray Structure  
of 3,9-[(H)<sub>2</sub>{P(p-tolyl)}<sub>3</sub>Ir]-3,9-μ-(H)<sub>2</sub>-7,8-C<sub>2</sub>B<sub>9</sub>H<sub>10</sub>.**

By

**10 James A. Doi, Raymond G. Teller, and M. Frederick Hawthorne\***

Prepared for Publication  
in

Chemical Communications

**DTIC  
ELECTE  
MAY 29 1980  
S D C**

Department of Chemistry  
University of California  
Los Angeles, California 90024

**14 12 May 12 1980**

Reproduction in whole or part is permitted  
for any purpose of the United States Government

Approved for Public Release; Distribution Unlimited

**DDC FILE COPY**

**30 5 21 018**

Reaction of  $[\text{Ir}(\text{cyclo-octa-1,5-diene})(\text{P}(\text{p-tolyl})_3)_2][\text{nido-7,8-C}_2\text{B}_9\text{H}_{12}]$   
with Hydrogen; Synthesis and X-ray Structure of  
 $3,9-\{(\text{H})_2(\text{P}(\text{p-tolyl})_3)_2\text{Ir}\}-3,9-\mu-(\text{H})_2-7,8-\text{C}_2\text{B}_9\text{H}_{10}$

By

James A. Doi, Raymond G. Teller and M. Frederick Hawthorne\*

Department of Chemistry  
University of California  
Los Angeles, California 90024

# SUMMARY

Treatment of the ion pair,  $[\text{Ir}(\text{cod})(\text{PR}_3)_2][\text{nido-7,8-C}_2\text{B}_9\text{H}_{12}]$  (cod = cyclo-octa-1,5-diene; R = phenyl, p-tolyl) with hydrogen gave closo-3,3-( $\text{PR}_3$ )<sub>2</sub>-3-H-3,1,2- $\text{IrC}_2\text{B}_9\text{H}_{11}$  and the novel nido-metallo-carborane, 3,9-[cis-(H)<sub>2</sub>-trans-( $\text{PR}_3$ )<sub>2</sub>-Ir]-3,9- $\mu$ -(H)<sub>2</sub>-nido-7,8-C<sub>2</sub>B<sub>9</sub>H<sub>10</sub>.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/_____	
Availability Codes	
Dist	Avail and/or special
A	

During our continuing investigation of the chemistry of the known alkene hydrogenation and isomerization catalyst; closo-3-3-(PPh<sub>3</sub>)<sub>2</sub>-3-H-3,1,2-RhC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>, we undertook the preparation of the iridium congener; closo-3-3-(PPh<sub>3</sub>)<sub>2</sub>-3-H-3,1,2-IrC<sub>2</sub>B<sub>9</sub>H<sub>11</sub> (Ia). Preparation of Ia is conveniently effected by treatment of [Ir(cod)(PPh<sub>3</sub>)<sub>2</sub>][nido-7,8-C<sub>2</sub>B<sub>9</sub>H<sub>12</sub>] with 1 atm. of hydrogen in cyclohexane heated to reflux. Characterization of Ia via nmr, infrared and elemental analysis was unequivocal.<sup>1</sup>

When the same reaction is carried out at -22°C in pentane, a mixture of products is formed. In addition to Ia, a pale yellow crystalline complex (IIa) is found in 40% yield. Infrared, <sup>1</sup>H, <sup>31</sup>P and <sup>11</sup>B nmr and microanalysis are consistent with the formulation; (H)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>Ir(C<sub>2</sub>B<sub>9</sub>H<sub>12</sub>). In particular, the <sup>1</sup>H nmr displays an envelope of broad signals between τ 13 ppm and τ 18 ppm, assigned to bridging Ir-H-B and B-H-B protons and high field signals at τ 28.0 and τ 28.5 ppm which are assigned to terminal metal hydrides. In addition, the infrared spectrum shows absorptions consistent with bridging Ir-H-B groups (ν 1965vw cm<sup>-1</sup>) and terminal Ir-H groups (ν 2249w cm<sup>-1</sup>).

Since a myriad of linkages between the transition metal and carborane cluster is possible, an X-ray structure analysis of the related (H)<sub>2</sub>(P(p-tolyl))<sub>2</sub>Ir-(C<sub>2</sub>B<sub>9</sub>H<sub>12</sub>) (IIb) was carried out.

Crystal Data: Ir[P(C<sub>7</sub>H<sub>7</sub>)<sub>3</sub>]<sub>2</sub>(H)<sub>2</sub>(C<sub>2</sub>B<sub>9</sub>H<sub>12</sub>)·C<sub>7</sub>H<sub>8</sub> M = 1028.5, triclinic, space group PT, a = 12.292(4), b = 12.854(3), c = 16.169(4)Å, α = 78.41(1)<sup>1</sup>, β = 94.88(1)<sup>0</sup>, γ = 102.04(1)<sup>0</sup>, Z = 2, ρ(calc 119K) = 1.40 g/cm<sup>3</sup>, ρ(obs) = 1.32 g/cm<sup>3</sup>, μ = (MoKα) = 31.3 cm<sup>-1</sup>. Data were collected on a Picker FACS-I automated diffractometer at -154°C. The structure was solved by standard Patterson-Fourier techniques. Full matrix least squares refinements of all atoms<sup>2</sup> (including the hydrogens bridging the Ir-B bonds) has converged to an agreement factor of 0.044.<sup>3</sup>

The molecule is illustrated in Figure 1, with a summation of bond distances and angles. The crystal structure formally consists of a discrete cis- $[(H)_2[(p\text{-tolyl})_3P]_2Ir]^+$  fragment bonded to two terminal B-H bonds of a  $[C_2B_9H_{12}]^{-1}$  anion.

---

Figure 1

---

The crystal displays an unusual disorder featuring the carborane cluster. Packing within the crystal is apparently determined by the tri-*p*-tolyl-phosphine ligands, leaving the carborane cluster free to adopt 1 of 2 different orientations in a 60:40 ratio. Each orientation of the carborane cluster however is constrained so that the Ir atom is bound to the same two B-H bonds; additionally the carbon atom positions within the carborane are also constrained to be identical in the two orientations. Consequently, the two species present in the crystal are chemically equivalent.

An especially noteworthy feature of the complex is the mode of interaction of the  $[(H)_2L_2Ir]^+$  fragment with the  $[C_2B_9H_{12}]^-$  moiety. The iridium atom interacts with the B(3)-H(3) and B(9)-H(9) bonds thereby completing the octahedral coordination about the metal atom. Exopolyhedral bonding of a transition metal to a borane cluster has been previously observed in  $[(Ph_3P)_2Cu]_2B_{10}H_{10} \cdot CHCl_3$ ,<sup>4</sup> (Cu-H = 1.97 (6) Å, Cu-B = 2.30(1) Å).

Comparison of Ir-B and Ir-H distances in IIb with those in  $Ir(B_5H_8)(CO)(PPh_3)_2$  (Ir-B (H bridged) = 2.250(6) Å, Ir-H = 1.73 Å),<sup>5</sup> a *nido*-pentagonal pyramid metalloborane, indicate a weaker Ir-carborane interaction in IIb (Ir-B = 2.46(1) Å, Ir-H = 1.94(8) Å). Note that this Ir-H bond length is also longer than the Ir-H bridging bond lengths found in the recently reported

$[(C_5Me_5)_2Ir_2(\mu-H)_3]BF_4$  (Ir-H 1.75(1) Å, neutron diffraction).<sup>6</sup>

Although the terminal Ir-H hydride ligands of IIb were not located in the structure analysis the geometry about the Ir atom is clearly that of a distorted octahedron with the symmetric hydrides occupying the two cis "vacant" positions.

Three center M-H-B (M = transition metal) bonding of exopolyhedral B-H units is seen in a variety of metallocarborane complexes.<sup>7</sup> However, all of these systems involve closo carboranyl or metallocarboranyl moieties. Previous to this work, the only M-H-B bonding of exopolyhedral B-H units of a nido carboranyl moiety were found in the main group compounds; 7,8- $\mu$ -dialkylaluminum-1,2-dicarba-nido-undecaborane (13) (alkyl = CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>) and its gallium analogues<sup>8,9</sup> in which the M-H-B bonds can be viewed as formally donating two electrons each to the aluminum or gallium to satisfy its "octet". The same formalism allows IIa to be seen as a stable eighteen electron organometallic compound.

---

Figure 2

---

Heating the nido compounds IIa or IIb, in benzene, gave the thermodynamically favored closo compounds Ia and its tri-*p*-tolylphosphine analogue Ib (both in 90% yield). This behavior is analogous to that of the aluminum and gallium nido-carborane systems.<sup>8</sup>

As previously noted, heating  $[Ir(cod)(PPh_3)_2][nido-7,8-C_2B_9H_{12}]$  in cyclohexane in the presence of hydrogen gave primarily the closo compound Ia. However, trace amounts of the nido compound IIa could also be isolated under those reaction conditions. Heating the  $[Ir(cod)(PR_3)_2][nido-7,8-C_2B_9H_{12}]$  (R=phenyl and *p*-tolyl) compounds in cyclohexane without hydrogen, gave only the closo compounds Ia and Ib, with no trace of the nido compounds.

Acknowledgements

This work was supported in part by the National Science Foundation (Grant No. CHE78-05679) and by the Office of Naval Research. We also wish to thank the National Science Foundation for the purchase of the Picker Diffractometer.




## References


1. a) T. E. Paxson and M. F. Hawthorne; J. Am. Chem. Soc., 96, 4674 (1974).  
b) T. E. Paxson, Ph.D. Thesis, University of California, Los Angeles (1974).
2. The p-tolyl rings were refined as rigid groups. After the location and refinement of all atoms except the four hydrogens bound to the Ir atom and the two partially occupied carborane hydrogens a difference-Fourier synthesis was calculated. The two tallest peaks in the map were attributed to the bridging hydrogen atoms. Their final refined positions are less than two standard deviations away from their Fourier map positions. The only other significant peak in the map in the vicinity of the Ir atom was in the proper position for a terminal hydride atom. Since the second hydride atom was not located (even in maps of decreasing  $\sin\theta/\tau$  cutoff) this atom was not included in the final cycles of refinement. The successful refinement of these bridging hydrogen atoms is due at least in part to their distance from the metal atom.
3. The crystal used for data collection was approximately equidimensional (0.10mm x 0.12mm x 0.14mm). An analytical absorption correction was made, the minimum and maximum absorption coefficients were 0.7462 and 0.8407 respectively. Structure factors were weighted by  $1/\sigma^2$  where  $\sigma$  is the standard deviation of F. In the final cycle of least squares no shift was larger than one-tenth of its corresponding estimated standard deviation.
4. J. T. Gill and S. J. Lippard; Inorg. Chem., 14, 751 (1974).
5. N. N. Greenwood, J. D. Kennedy, W. S. McDonald, D. Reed and J. Staves; J.C.S. Dalton, 117 (1979).
6. R. Bau, W. E. Carroll, D. W. Hart, R. G. Teller and T. F. Koetzle; Adv. Chem. Ser., 167, 73 (1978).
7. a) R. T. Baker, R. E. King III, C. Knobler, C. A. O'Con and M. F. Hawthorne; J. Am. Chem. Soc., 100, 8266 (1978).  
b) G. Allegra, M. Calligaris, R. Furlanetto, G. Nardin, and L. Randaccio; Cryst. Struct. Commun., 3, 69 (1974).  
c) R. A. Love and R. Bau; J. Am. Chem. Soc., 94, 8274 (1972).
8. D. A. T. Young, R. J. Wiersema and M. F. Hawthorne; J. Am. Chem. Soc., 93, 5687 (1971).

9. a) The cage numbering system in this paper is consistent with that favored by IUPAC for nido carboranes, and is different from the numbering scheme in ref. 6.

b) Pure Appl. Chem.; 30, 683 (1972).

Figure 1 The molecular structure of  $3,9-\{(\text{H})_2(\text{P}(\text{p-tolyl})_3)_2\text{Ir}\}-3,9-\mu-(\text{H})_2-7,8-\text{C}_2\text{B}_9\text{H}_{10}$ . Some important bond lengths are: Ir-P(1), 2.312(3); Ir-P(2), 2.301(3); Ir-H(3), 1.85(7); Ir-H(9), 2.02(10); Ir-B(3), 2.480(11); Ir-B(9), 2.452(11); B(3)-H(3), 1.4(1); B(9)-H(9), 1.3(1) Å; bond angles: P(1)-Ir-P(2), 158.9(1); P(1)-Ir-H(3), 102(2); P(1)-Ir-H(9), 94(3); P(2)-Ir-H(3), 94(2); P(2)-Ir-H(9), 95(3); H(3)-Ir-H(9), 104(4); B(3)-Ir-B(9), 40.8(4); Ir-H(3)-B(3), 98(4); Ir-H(9)-B(9), 94(6)°.





Cod = cyclo-octa-1,5-diene

a) R = Ph

b) R = p-tolyl

Figure 2

Some Distances (in Å) and Angles (in degrees) in 3,9- $\{(\text{H})_2(\text{P}(\text{p-tolyl})_3)_2\text{Ir}\}$ -  
3,9- $\mu\text{-(H)}_2\text{-7,8-C}_2\text{B}_9\text{H}_{10}$ .

Distances about the Ir and P atoms

Ir-P(1)	2.312(3)	P(1)-Group 1	1.807
Ir-P(2)	2.301(3)	P(1)-Group 2	1.828
Ir-H(3)	1.85(7)	P(1)-Group 3	1.829
Ir-H(9)	2.02(10)	P(2)-Group 4	1.831
Ir-B(3)	2.480(11)	P(2)-Group 5	1.814
Ir-B(9)	2.452(11)	P(2)-Group 6	1.859

Distances within the carborane fragment<sup>a</sup>

B(1)-B(2)	1.56(3)	B(5)-B(6)	1.68(3)
B(1)-B(3)	1.73(2)	B(5)-B(10)	1.70(3)
B(1)-B(4)	1.72(2)	B(5)-B(11)	1.71(3)
B(1)-B(5)	1.81(3)	B(6)-C(7)	1.70(2)
B(1)-B(6)	1.79(3)	B(6)-B(11)	1.77(3)
B(2)-B(3)	1.77(2)	C(7)-C(8)	1.58(1)
B(2)-B(6)	1.60(3)	C(7)-B(11)	1.67(2)
B(2)-C(7)	1.73(3)	C(7)-B(2)'	1.71(3)
B(2)-C(8)	1.73(3)	C(8)-B(9)	1.72(2)
B(3)-B(4)	1.71(2)	C(8)-B(2)'	1.73(3)
B(3)-C(8)	1.69(2)	B(9)-B(10)	1.78(2)
B(3)-B(9)	1.72(2)	B(9)-B(2)'	1.76(2)
B(4)-B(5)	1.67(2)	B(10)-B(11)	1.75(2)
B(4)-B(9)	1.72(2)	B(10)-B(2)'	1.69(3)
B(4)-B(10)	1.72(2)	B(11)-B(2)'	1.63(3)
B(1)-H(1)	1.3(2)	C(7)-H(7)	1.0(3)
B(3)-H(3)	1.4(1)	C(8)-H(8)	1.2(1)
B(4)-H(4)	1.2(2)	B(9)-H(9)	1.3(1)
B(5)-H(5)	0.9(1)	B(10)-H(10)	1.2(1)
B(6)-H(6)	1.0(1)	B(11)-H(11)	1.0(2)

Angles

P(1)-Ir-P(2)	158.9(1)	H(3)-Ir-H(9)	104(4)
P(1)-Ir-H(3)	102(2)	B(3)-Ir-B(9)	40.8(4)
P(1)-Ir-H(9)	94(3)	Ir-H(3)-B(3)	98(4)
P(2)-Ir-H(3)	94(2)	Ir-H(9)-B(9)	94(6)
P(2)-Ir-H(9)	95(3)		

- a) A numbering system consistent with that of figure 1 is used. B(2) and B(2)' are one half weighted atoms.

1. Positional (in fractional coordinates) and Thermal Parameters<sup>a</sup> for the Nongroup Atoms of 3,9-[(H)<sub>2</sub>(P(p-tolyl)<sub>3</sub>)<sub>2</sub>Ir]-3,9-μ-(H)<sub>2</sub>-7,8-C<sub>2</sub>B<sub>9</sub>H<sub>10</sub>.

Atom <sup>b</sup>	<u>x</u>	<u>y</u>	<u>z</u>	<u>B(A)<sup>02</sup></u>
Ir	.17641(3)	.40184(3)	.32659(3)	c
P(1)	.2165(2)	.5270(2)	.2034(2)	c
P(2)	.1783(2)	.3285(2)	.4686(2)	c
B(1)[B(10)]	.0291(16)	.1891(15)	.1765(13)	5.5(4)
B(2)[---]	-.0043(19)	.1140(18)	.2624(15)	1.6
B(3)[B(9)]	.0743(9)	.2457(9)	.2646(7)	1.7(2)
B(4)[B(4)]	.0101(12)	.3170(12)	.1805(9)	3.2(3)
B(5)[B(5)]	-.0985(16)	.2369(16)	.1418(14)	5.2(4)
B(6)[B(11)]	-.1074(17)	.1088(18)	.1950(13)	5.6(4)
C(7)[C(7)]	-.1344(12)	.1169(12)	.2944(9)	4.4(3)
C(8)[C(8)]	-.0311(10)	.1924(10)	.3309(8)	3.3(2)
B(9)[B(3)]	-.0144(9)	.3259(9)	.2820(7)	1.5(2)
B(10)[B(1)]	-.1223(15)	.3244(15)	.2016(11)	4.5(4)
B(11)[B(6)]	-.1974(15)	.1910(14)	.2155(11)	4.5(4)
---[B(2)]	-.1489(19)	.2455(18)	.2972(15)	1.6
H(1)[H(10)]	.081(9)	.175(9)	.111(7)	4.(3)
H(3)[H(9)]	.184(6)	.273(6)	.293(5)	1.(2)
H(4)[H(4)]	.060(12)	.391(12)	.131(10)	8.(4)
H(5)[H(5)]	-.110(9)	.267(9)	.090(7)	2.(3)
H(6)[H(11)]	-.119(8)	.032(9)	.211(6)	2.(2)
H(7)[H(7)]	-.196(21)	.060(20)	.320(16)	13.(8)
H(8)[H(8)]	-.010(7)	.159(6)	.403(5)	1.(2)
H(9)[H(3)]	.012(9)	.405(9)	.318(7)	3.(3)
H(10)[H(1)]	-.160(8)	.396(8)	.165(6)	2.(2)
H(11)[H(6)]	-.277(18)	.168(17)	.217(13)	16.(7)

2. Anisotropic Temperature Factors<sup>d</sup>

Atom	<u>10<sup>5</sup>β<sub>11</sub></u>	<u>10<sup>5</sup>β<sub>22</sub></u>	<u>10<sup>5</sup>β<sub>33</sub></u>	<u>10<sup>5</sup>β<sub>12</sub></u>	<u>10<sup>5</sup>β<sub>13</sub></u>	<u>10<sup>5</sup>β<sub>23</sub></u>
Ir	228(3)	206(3)	82(2)	34(2)	23(2)	-14(1)
P(1)	277(20)	213(18)	144(11)	47(15)	64(12)	-17(11)
P(2)	294(20)	266(18)	97(11)	36(15)	15(12)	-41(11)



### 3. Rigid Group Parameters<sup>e</sup>

<u>Group</u>	<u>x</u>	<u>y</u>	<u>z</u>	<u>Phi</u>	<u>Theta</u>	<u>Rho</u>	<u>B(A<sup>2</sup>)</u>
Ptoly1 1	.2529(5)	.2154(4)	.5000(4)	2.436(3)	2.614(4)	2.783(3)	2.14(7)
Ptoly1 2	.0446(4)	.2833(5)	.5189(4)	-2.642(3)	-2.561(4)	0.504(4)	2.28(9)
Ptoly1 3	.2516(4)	.4290(4)	.5336(4)	1.917(5)	-2.382(3)	0.925(4)	1.87(7)
Ptoly1 4	.1013(4)	.5966(4)	.1689(4)	-3.138(7)	2.042(4)	-.586(6)	2.07(7)
Ptoly1 5	.2633(5)	.4906(4)	.1101(3)	0.234(4)	2.585(3)	-1.122(3)	1.84(7)
Ptoly1 6	.3347(4)	.6376(4)	.2172(4)	-2.351(3)	-2.556(4)	3.128(3)	2.00(7)
Ptoly1 7 <sup>f</sup>	.4837(8)	.1341(13)	.1055(9)	.54(2)	-1.95(1)	3.18(2)	2.8(2)
Ptoly1 8 <sup>f</sup>	.4679(13)	.1412(17)	.1539(13)	-1.80(2)	2.13(1)	-.75(2)	5.0(3)

- Estimated standard deviations in the least significant figures are given in parentheses.
- Alternate atomic designations for the carborane cage atoms are given in brackets. Note that B(2) is a partially weighted atom.
- Anisotropic thermal parameters.
- The form of the anisotropic thermal ellipsoidal is:  

$$\exp[-(\beta_{11}h^2 + \beta_{22}k^2 + \beta_{33}l^2 + 2\beta_{12}hk + 2\beta_{13}hl + 2\beta_{23}kl)]$$
- x, y, and z are the fractional coordinates of the origin of the rigid group and phi, theta, and rho the rotation angles (in radians)
- One half weighted groups.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report 109	2. GOVT ACCESSION NO. AD-A084 837	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Reaction of [Ir(cyclo-octa-1,5-diene) (P(p-tolyl) <sub>3</sub> ) <sub>2</sub> ][nido-7,8-C <sub>2</sub> B <sub>9</sub> H <sub>12</sub> ] with Hydrogen; Synthesis and X-Ray Structure of 3,9-[(H) <sub>2</sub> {P(p- tolyl) <sub>3</sub> } <sub>2</sub> Ir]-3,9-μ-(H) <sub>2</sub> -7,8-C <sub>2</sub> B <sub>9</sub> H <sub>10</sub>		5. TYPE OF REPORT & PERIOD COVERED Interim
7. AUTHOR(s) James A. Doi, Raymond G. Teller and M.F. Hawthorne		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS The University of California Department of Chemistry Los Angeles, California 90024		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0390
11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Branch Office of Naval Research Washington, D.C. 20360		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-608
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE May 12, 1980
		13. NUMBER OF PAGES 12 incl. figures
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Metal Hydride Bridges Iridium Complex Metallocarborane Metal Cluster		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Treatment of the ion pair, [Ir(cod)(PR <sub>3</sub> ) <sub>2</sub> ][nido-7,8-C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> ] (cod=cyclo-octa- 1,5-diene; R=phenyl, p-tolyl) with hydrogen gave closb-3,3-(PR <sub>3</sub> ) <sub>2</sub> -3-H-3,1,2- IrC <sub>2</sub> B <sub>9</sub> H <sub>11</sub> and the novel nido-metallocarborane, 3,9-[cis-(H) <sub>2</sub> -trans-(PR <sub>3</sub> ) <sub>2</sub> -Ir]- 3,9-μ-(H) <sub>2</sub> -nido-7,8-C <sub>2</sub> B <sub>9</sub> H <sub>10</sub> .		

DD FORM 1 JAN 75 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

APPENDIX

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. R. N. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1	Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1
Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1	Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1
		Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1	Professor O. T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214	1
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1	Professor P. S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1	Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1	Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
Professor H. Abrahamson University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1	Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Branch Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	
ONR Branch Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.  
Copies

Dr. Rudolph J. Marcus  
Office of Naval Research  
Scientific Liaison Group  
American Embassy  
APO San Francisco 96503

1

Mr. James Kelley  
DTNSRDC Code 2803  
Annapolis, Maryland 21402

1